

BRIEF REPORT

WILEY

Dynamic ^{18}F -FDG PET/CT can predict the major pathological response to neoadjuvant immunotherapy in non-small cell lung cancer

Zhi-Yong Chen¹ | Rui Fu^{2,3} | Xiao-Yue Tan⁴ | Li-Xu Yan⁵ | Wen-Fang Tang⁶  |
Zhen-Bin Qiu³ | Yi-Fan Qi^{2,3} | Yu-Fa Li⁵ | Qing-Yi Hou⁴ | Yi-Long Wu^{1,2,3} |
Wen-Zhao Zhong^{1,2,3}  | Ben-Yuan Jiang¹

¹Guangdong Lung Cancer Institute, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, China

²School of Medicine, South China University of Technology, Guangzhou, China

³Guangdong Provincial Key Laboratory of Translational Medicine in Lung Cancer, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, China

⁴Department of Nuclear Medicine, WeiLun PET/CT Center, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, China

⁵Department of Pathology, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, China

⁶Department of Cardiothoracic Surgery, Zhongshan City People's Hospital, Zhongshan, Guangdong, China

Correspondence

Wen-Zhao Zhong, Guangdong Lung Cancer Institute, Guangdong Provincial Key Laboratory of Translational Medicine in Lung Cancer, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, China; School of Medicine, South China University of Technology, Guangzhou, China.
Email: 13609777314@163.com

Ben-Yuan Jiang, Guangdong Lung Cancer Institute, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, China.
Email: jiangben1000@126.com

Funding information

GDPH Scientific Research Funds for Leading Medical Talents and Distinguished Young Scholars in Guangdong Province, Grant/Award Number: KJ012019449; Guangdong Basic and Applied Basic Research Foundation, Grant/Award Number: 2019B1515130002; Guangdong Provincial People's Hospital Young Talent Project, Grant/Award Number: GDPPHYTP201902; High-level Hospital Construction Project, Grant/Award Numbers: DFJH201801, DFJH201910; National Natural Science Foundation of China, Grant/Award Numbers: 81673031, 81872510; Research Fund from Guangzhou Science and Technology Bureau, Grant/Award Number: 201704020161

Abstract

Major pathological response (MPR) is a potential surrogate for overall survival. We determined whether the dynamic changes in ^{18}F -labeled fluoro-2-deoxyglucose positron emission tomography/computed tomography (^{18}F -FDG PET/CT) were associated with MPR in patients receiving neoadjuvant immunotherapy. Forty-four patients with stage II–III non-small cell lung cancer (NSCLC) who received neoadjuvant immunotherapy and radical surgery were enrolled. Moreover, ^{18}F -FDG PET/CT scans were performed at baseline and within 1 week before surgery to evaluate the disease. All histological sections were reviewed to assess MPR. The detailed clinical features of the patients were analyzed. The reliability of the clinical variables was assessed in differentiating between MPR and non-MPR using logistic regression. Receiver-operating characteristic (ROC) curve analysis identified the SUVmax changes threshold most associated with MPR. Most of the patients were pathologically diagnosed with squamous cell carcinoma and received anti-PD-1 antibodies plus chemotherapy. The immunotherapy regimens included nivolumab, pembrolizumab, and camrelizumab. MPR was observed in more than half of lesions. Tumors with MPR had a higher decrease in the longest dimension on dynamic PET/CT than those without MPR. Furthermore, the decline in SUVmax was significantly different between MPR and non-MPR diseases, and MPR lesions had a prominent mean reduction in SUVmax. SUVmax reduction was independently associated with MPR in the multivariate regression. On ROC analysis, the threshold of SUVmax decrease in 60% was associated with MPR. Dynamic changes in SUVmax were associated with MPR. The tumors with MPR showed a greater PET/CT response than those without MPR. A SUVmax decrease of more than 60% is more likely to result in an MPR after receiving neoadjuvant immunotherapy.

Zhi-Yong Chen, Rui Fu, Xiao-Yue Tan, and Li-Xu Yan contributed equally to this work.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Thoracic Cancer* published by China Lung Oncology Group and John Wiley & Sons Australia, Ltd.

KEYWORDS¹⁸F-FDG PET/CT, immunotherapy, major pathological response, neoadjuvant therapy, non-small cell lung cancer**INTRODUCTION**

Stage II–III non-small cell lung cancer (NSCLC) has high heterogeneity, involves multidisciplinary and different therapy regimens, and shows distinct prognoses. Patients with stage II–III disease still have a high risk of recurrence, even if they undergo radical surgery.¹ Preoperative chemotherapy is recommended for patients with resectable NSCLC, but it has limited survival benefits.^{1–3} Although data with a longer follow-up duration are not mature, immunotherapy as a neoadjuvant treatment shows encouraging efficacy.^{4–6} Major pathological response (MPR) has been used in several randomized clinical trials as an exploratory endpoint and is expected to be associated with survival.^{5–8}

¹⁸F-labeled fluoro-2-deoxyglucose positron emission tomography/computed tomography (¹⁸F-FDG PET/CT) is a helpful tool that can reflect the tumor size and glycometabolism in the whole body and is widely used in evaluating tumor stages and therapeutic responses. It is important to determine patients who have a great response to immunotherapy because incorrect evaluation judgment may lead to making incorrect decisions about the next step of treatment after induced immunotherapy. In particular, discordance occurred between the CT Response Evaluation Criteria in Solid Tumors and the histopathological response.⁹ Progression or stable CT at a single time does not mean that the disease has a poor response to immunotherapy or has no potential opportunity to receive surgery. Some researchers have suggested that ¹⁸F-FDG PET/CT may be more accurate in predicting pathological responses to neoadjuvant chemotherapy.^{10,11} Nivolumab,¹² pembrolizumab,¹³ and camrelizumab¹⁴ have shown great effect in locally advanced NSCLC. In this real-world retrospective study, we performed analyses to determine whether the dynamic changes in ¹⁸F-FDG PET/CT were associated with MPR in patients receiving different preoperative immunotherapies.

METHODS**Patients**

We retrospectively enrolled patients with stage II–III (eighth edition) NSCLC at baseline who had received preoperative immune checkpoint inhibitors treatment and underwent surgery at Guangdong Provincial People's Hospital between July 2019 and July 2021. ¹⁸F-FDG PET/CT scans were performed at baseline and within 1 week before surgery. All primary tumors and lymph nodes were subjected to standard MPR evaluation by two pathologists. Detailed clinicopathological characteristics were collected for analysis. This study was approved by the research ethics committee of the

hospital (GDREC2016175H[R2]). Written informed consent was obtained from all the patients.

Major pathological response assessments

All histological sections were reviewed by two pathologists following the guidelines of the International Association for the Study of Lung Cancer (IASLC) Multidisciplinary Recommendations for Pathologic Assessment of Lung Cancer Resection Specimens After Neoadjuvant Therapy.¹⁵ The primary components of the tumor bed, including viable tumor, necrosis, and stroma were described and recorded. MPR is defined as ≤10% of the viable tumor.

TABLE 1 Demographics of the cohort

Patient characteristics		N (%)
		44 (100%)
Age, years, mean (IQR)		59.2 (53–67)
Sex (n,%)		
	Female	7 (15.9%)
	Male	37 (84.1%)
Smoker		
	Yes	30 (68.2%)
	No	14 (31.8%)
Pathological diagnosis		
	LUAD	9 (20.4%)
	SCC	27 (61.4%)
	LELC	8 (18.2%)
TNM stage (eighth)		
	II	9 (20.5%)
	III	35 (79.6%)
PD-L1, mean (IQR)		32% (1%–60%)
Neoadjuvant regimens		
	Immunotherapy	3 (6.8%)
	Immunotherapy plus chemotherapy	41 (93.2%)
Immunotherapy regimens		
	Nivolumab	29 (65.9%)
	Pembrolizumab	8 (18.2%)
	Camrelizumab	7 (15.9%)
Chemotherapy regimens (if applicable)		
	Paclitaxel + platinum	36 (81.8%)
	Pemetrexed + platinum	5 (11.4%)

¹⁸F-labeled fluoro-2-deoxyglucose positron emission tomography/computed tomography scans

¹⁸F-FDG PET/CT scans were performed using Siemens Biograph16 when treatment-naïve and within 1 week before surgery. Before the scan, all patients needed to fast for more than 4–6 h and were required to have blood glucose levels less than 7.0 mmol/l. The dose of ¹⁸F-FDG was 0.16 mCi/Kg. PET/CT

images were obtained in 60–70 min after ¹⁸F-FDG injection. From head to thigh, the scan speeds were 5 min/bed (head, 1 bed) and 2 min/bed (skull base to thigh, 5–8 beds). Noncontrast CT was used to adjust the attenuation with a tube voltage of 120 kV and a tube current of 50 mAs. The tube voltage of the spiral contrast-enhanced CT was 120 kV, and the tube currents were 300 mAs (head), 140 mAs (chest), and 180 mAs (abdomen). The images were independently assessed and evaluated by two nuclear medicine specialists.




FIGURE 1 A 41-year-old man was diagnosed with squamous cell carcinoma and received three cycles of nivolumab plus paclitaxel (albumin-bound) and carboplatin. (a) Positron emission tomography/computed tomography (PET/CT) scans at baseline, (b) PET/CT scans within 1 week before surgery




FIGURE 2 A 48-year-old woman was diagnosed with lymphoepithelioma-like carcinoma and received three cycles of pembrolizumab plus pemetrexed and carboplatin. (a) Positron emission tomography/computed tomography (PET/CT) scans at baseline, (b) PET/CT scans within 1 week before surgery

FIGURE 3 Hematoxylin and eosin staining showed that the patient from Figure 1 had a major pathological response (MPR) and the residual viable tumor was less than 10%, and the residual viable tumor was more than 10% in the patient from Figure 2. (a) MPR with 40× and (b) 100×; (c) non-MPR with 40× and (d) 100×




TABLE 2 Clinical characteristics

Characteristics	Tumor with MPR (<i>n</i> = 28)	Tumor without MPR (<i>n</i> = 16)	<i>p</i> -value
Age, years, mean (IQR)	59.5 (53–67)	57.9 (54–67)	0.55 ^a
Male	25 (89.3%)	12 (75.0%)	0.41 ^b
Smoker	19 (67.9%)	11 (68.8%)	0.95 ^c
Pathological diagnosis			0.06 ^d
LUAD	4 (14.3%)	5 (31.3%)	
SCC	21 (75.0%)	6 (37.5%)	
LELC	3 (10.7%)	5 (31.3%)	
TNM stage (eighth)			0.55 ^b
II	7 (25.0%)	2 (12.5%)	
III	21 (75.0%)	14 (87.5%)	
PD-L1, mean (IQR)	36.3% (1%–80%)	25.3% (5%–33%)	0.31 ^a
Neoadjuvant regimens			1 ^b
Immunotherapy	2 (7.1%)	1 (6.3%)	
Immunotherapy plus chemotherapy	26 (92.9%)	15 (93.8%)	
Immunotherapy regimens			0.22 ^d
Nivolumab	21 (75.0%)	8 (50.0%)	
Pembrolizumab	4 (14.3%)	4 (25.0%)	
Camrelizumab	3 (10.7%)	4 (25.0%)	
Chemotherapy regimens (if applicable)			0.10 ^b
Paclitaxel ^e + platinum	25 (96.2%)	11 (73.3%)	
Pemetrexed + platinum	1 (4.8%)	4 (26.7%)	

^aAccording to a Student's *t*-test.

^bAccording to a correction for the continuity χ^2 test.

^cAccording to a Pearson's χ^2 test.

^dAccording to a Fisher's precision test.

^ePaclitaxel (albumin-bound).

Abbreviations: LELC, lymphoepithelioma-like carcinoma; LUAD, lung adenocarcinoma; SCC, squamous cell carcinoma.

Statistical analysis

Consecutive data are presented as median $\pm$ standard deviation or interquartile range (IQR). Student's *t*-tests were used to compare differences in the parameter variables. Categorical variables were analyzed using the correction for continuity χ^2 test, Pearson χ^2 test, or Fisher's precision test. Logistic regression was performed to determine the independent association between clinical variables and MPR. Subsequently, the sensitivity and specificity were explored via receiver-operating characteristic (ROC) curve analysis to determine the dynamic SUVmax changes threshold that could best separate MPR from non-MPR. All statistical analyses were performed using the International Business Machines (IBM) Statistical Package for the Social Sciences Statistics version 26.0 (IBM Corp.). Statistical significance was set at $p < 0.05$.

TABLE 3 Analyses of dynamic positron emission tomography/computed tomography scans

Characteristics	Tumor with MPR	Tumor without MPR	<i>p</i> -value
Longest dimension (cm)			
At baseline	5.3 $\pm$ 3.5	5.1 $\pm$ 1.8	0.85 ^a
Before surgery	2.7 $\pm$ 2.0	3.8 $\pm$ 1.5	0.07 ^a
Changing	−45.1% $\pm$ 30.5%	−25.1% $\pm$ 22.9%	<0.05 ^a
SUVmax			
At baseline	14.8 $\pm$ 5.0	15.7 $\pm$ 5.2	0.58 ^a
Before surgery	2.6 $\pm$ 1.6	8.6 $\pm$ 6.5	<0.05 ^a
Changing	−80.4% $\pm$ 13.7%	−46.5% $\pm$ 33.0%	<0.05 ^a

^aAccording to a Student's *t*-test.

TABLE 4 Logistic regression for major pathological response

Variables	<i>p</i> -value	OR	95% CI
PD-L1	0.15	0.14	0.01–2.05
Size reduction	0.508	3.13	0.11–91.22
SUVmax reduction	0.01	386.45	4.14–36101.23

Abbreviations: CI, confidence interval; OR, odds ratio.

TABLE 5 Receiver-operator characteristic (ROC) threshold and for major pathological response

		MPR (<i>n</i> = 28)	Without MPR (<i>n</i> = 16)	Total	Sensitivity %	Specificity %	PPV %	NPV %
ROC threshold	SUVmax decreasing $\geq 60.0\%$	25	6	31	89.3	62.5	80.6	76.9
	SUVmax decreasing <60.0%	3	10	13				

Abbreviations: NPV, negative predictive value; PPV, positive predictive value.

RESULTS

A total of 44 patients were finally enrolled in this study (Table 1). Most were male (37/44, 84.1%) and smokers (30/44, 68.2%), and mean age was 59.2 (IQR, 53–67) years. Almost two-thirds of the tumors (28/44, 63.6%) showed MPR (Figure 1), and the rest were non-MPR (16/44, 36.4%) (Figure 2) Hematoxylin and eosin staining examples are shown in Figure 3. The majority of the pathological diagnoses were squamous cell carcinoma (SCC, 27/44, 61.4%), followed by adenocarcinoma (9/44, 20.5%), and lymphoepithelioma-like carcinoma (8/44, 18.2%). Almost 80% of the patients (35/44, 79.6%) had stage III NSCLC (eighth edition) at baseline, and most of them finally received three cycles of immunotherapy plus chemotherapy (41/44, 93.2%). Approximately 60% of the patients received nivolumab (29/44, 65.9%), 18% received pembrolizumab (8/44, 18.2%), and 16% received camrelizumab (7/44, 15.9%) as immunotherapy regimens. Paclitaxel (albumin-bound) and platinum (36/44, 81.8%) were the dominant chemotherapy regimens. There were no significant differences between tumors with MPR and without MPR in terms of age, sex, smoking history, pathological diagnosis, tumor-node-metastasis (TNM) stages, programmed death-ligand 1 (PD-L1), neoadjuvant, immunotherapy, and chemotherapy regimens (Table 2).

Further analyses showed that the longest dimension ($p = 0.85$) and SUVmax ($p = 0.58$) of ¹⁸F-FDG PET/CT at baseline were not significantly different between the tumor with MPR and without MPR. SUVmax before surgery showed a significant difference ($p < 0.05$), whereas the longest dimension did not ($p = 0.07$). The dynamic longest dimension ($p < 0.05$) and SUVmax ($p < 0.05$) demonstrated significant differences between the MPR and non-MPR groups and have a remarkable decline in two the groups (Table 3). This suggests that dynamic decreases in dimension and SUVmax value of the tumor are associated with tumor MPR.

Logistic regression was used to determine the association between variables and the prediction of MPR, as opposed to non-MPR. SUVmax reduction demonstrated an odds ratio of 386.45, with a 95% confidence interval of 4.14–36101.23 (p -value <0.05) (Table 4). ROC analysis showed that a dynamic SUVmax decrease of 60% was the ideal threshold. SUVmax decreasing $\geq 60.0\%$ was associated with MPR, with 0.83 area under the curve, 89.3% sensitivity, 62.5%




FIGURE 4 Receiver-operating characteristic curve of decreasing SUVmax used to differentiate major pathological response from nonmajor pathological response

specificity, 80.6% positive predictive value, and 76.9% negative predictive value (Table 5, Figure 4).

DISCUSSION

Preoperative immunotherapy has demonstrated interesting short-term outcomes in several neoadjuvant trials^{4–6} and is expected to become a standard treatment pending long-term efficacy releasing. MPR is considered as a vital short-term endpoint that has the potential to replace long-term survival outcomes in neoadjuvant immunotherapy. However, a discrepancy has been observed between the CT and pathological responses.⁹ Stromal, fibrous, inflammatory, and necrotic components may confuse the assessment of radiological response in tumor size and interfere with predicting pathological response after neoadjuvant immunotherapy and before surgery.¹⁵ Previous studies have demonstrated that modifications in metabolic activity, represented by changes in the SUV, are associated with tumor response.¹⁶ ¹⁸F-FDG PET/CT has showed its value for predicting pathological response in patients with NSCLC who underwent neoadjuvant chemotherapy.^{10,11} In this study, we explored the correlation between PET/CT and MPR.

In this real-world retrospective study, all patients with stage II–III NSCLC received three cycles of neoadjuvant therapy, radical surgery, and dynamic ¹⁸F-FDG PET/CT scans. Surgical specimens were reviewed to assess MPR. In terms of clinical characteristics, pathological diagnosis between the patients with and without MPR showed a marginal statistical difference, and the MPR groups had a higher proportion of SCC than the non-MPR group. In the real

world, locally advanced patients with SCC are more likely to received neoadjuvant immunotherapy than those with adenocarcinoma. This was probably because SCC has lower mutation rates of targeted genes than adenocarcinoma. Peri-operative targeted therapy has shown primary efficacy in patients with resectable adenocarcinoma.^{17,18} However, it remains unclear whether patients with sensitizing epidermal growth factor receptor mutations or anaplastic lymphoma kinase alterations would benefit from neoadjuvant immunotherapy. These patients were excluded from some neoadjuvant immunotherapy clinical trials, such as the CheckMate-816,⁶ probably because the patients with these targeted driving mutations had risks of hyperprogression and other adverse events related to immunotherapy,¹⁹ and had limited efficiency.²⁰

Response patterns of immunotherapy are different from those of targeted therapy or chemotherapy; sometimes, pseudoprogression and hyperprogression would disturb response assessment.²¹ FDG PET/CT plays a vital role in evaluating the response of solid tumors to immunotherapy. One of the main reasons for this is that the SUV is a parameter that shows the tumor metabolic activity. Immune infiltrates are associated with better immunotherapy responses. A significant correlation was demonstrated between the SUV and the expression of PD-L1²² and CD8-tumor-infiltrating-lymphocytes²³ at baseline. PD-L1-expressing NSCLC has a high glucose metabolism with a high SUVmax.²² Therefore, PET/CT has the potential to reflect some characteristics of the tumor immune microenvironment and predict the response to immunotherapy. Compared with a previous study that reported that metabolic parameters calculated by PET were significantly correlated with the pathological response in patients who received two cycles of sintilimab,²⁴ our cohort had more diverse therapeutic regimens and evaluated MPR under IASLC multidisciplinary recommendations. In this study, there was no significant difference between the MPR and non-MPR groups in other clinical characteristics, including TNM stage, PD-L1 expression, and therapy regimens. Tumors with MPR showed a greater decrease in longest dimension and SUVmax. We found that SUVmax reduction in tumors on dynamic PET/CT is strongly and independently associated with MPR and provides high accuracy. In our cohort, and SUVmax decrease of more than 60% was more likely to have an MPR. These results emphasized that dynamic ¹⁸F-FDG PET/CT is important in evaluating the clinical response of preoperative immunotherapy and can help to determine the population who can benefit from neoadjuvant immunotherapy followed by radical surgery.

This study had some limitations. First, it was retrospective with a limited small sample size. Further investigation is required to validate these results. Second, most patients received immunotherapy plus chemotherapy, with nivolumab + paclitaxel + platinum as a regimen. SCC is the most common pathological type. This may cause bias that interferes with the effectiveness of the results of this study, although it partly reflected the real-world situation of

neoadjuvant immunotherapy in East Asia. Third, overall survival data were not mature; therefore, we could not determine the association between ^{18}F -FDG PET/CT, MPR, and prognosis.

In conclusion, dynamic changes in SUVmax on ^{18}F -FDG PET/CT were associated with MPR in patients who received neoadjuvant immunotherapy. Tumors with MPR showed a greater ^{18}F -FDG PET/CT response than those without MPR. An SUVmax decrease of more than 60% is more likely to result in an MPR after receiving preoperative immunotherapy.

ACKNOWLEDGMENTS

This research was supported by funding from the National Natural Science Foundation of China (81673031, 81872510), Guangdong Provincial People's Hospital Young Talent Project (GDPHYTP201902), Guangdong Basic and Applied Basic Research Foundation (no. 2019B1515130002), High-level Hospital Construction Project (DFJH201801, DFJH201910), Research Fund from Guangzhou Science and Technology Bureau (no. 201704020161), GDPH Scientific Research Funds for Leading Medical Talents and Distinguished Young Scholars in Guangdong Province (no. KJ012019449).

CONFLICT OF INTEREST

YLW is a consultant of AstraZeneca, Roche Holdings AG, and Boehringer Ingelheim; he has received honoraria from AstraZeneca, Eli Lilly, Roche Holdings AG, Pfizer, Sanofi, Boehringer Ingelheim, Merck Sharp & Dohme, and Bristol-Myers Squibb; he has received research funding from Boehringer Ingelheim (Inst), Roche Holdings AG (Inst). WZZ has received honoraria from AstraZeneca, Eli Lilly, Roche Holdings AG, Pfizer, Sanofi, Boehringer Ingelheim, Merck Sharp & Dohme, and Bristol-Myers Squibb.

ORCID

Wen-Fang Tang  <https://orcid.org/0000-0001-6116-6838>

Wen-Zhao Zhong  <https://orcid.org/0000-0002-8917-8635>

REFERENCES

- Arriagada R, Bergman B, Dunant A, Le Chevalier T, Pignon JP, Vansteenkiste J. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small-cell lung cancer. *N Engl J Med*. 2004;350(4):351–60.
- Pignon JP, Tribodet H, Scagliotti GV, Douillard JY, Shepherd FA, Stephens RJ, et al. Lung adjuvant cisplatin evaluation: a pooled analysis by the LACE collaborative group. *J Clin Oncol*. 2008;26(21):3552–9.
- NSCLC Meta-analysis Collaborative Group. Preoperative chemotherapy for non-small-cell lung cancer: A systematic review and meta-analysis of individual participant data. *Lancet*. 2014;383(9928):1561–71.
- Forde PM, Chaft JE, Smith KN, Anagnostou V, Cottrell TR, Hellmann MD, et al. Neoadjuvant PD-1 blockade in Resectable lung cancer. *N Engl J Med* 2018;378(21):1976–86.
- Shu CA, Gainor JF, Awad MM, Chiuhan C, Grigg CM, Pabani A, et al. Neoadjuvant atezolizumab and chemotherapy in patients with resectable non-small-cell lung cancer: an open-label, multicentre, single-arm, phase 2 trial. *Lancet Oncol*. 2020;21(6):786–95.
- Forde PM, Spicer J, Lu S, Provencio M, Mitsudomi T, Awad MM, et al. Abstract CT003: Nivolumab (NIVO) + platinum-doublet chemotherapy (chemo) vs chemo as neoadjuvant treatment (tx) for resectable (IB-IIIa) non-small cell lung cancer (NSCLC) in the phase 3 CheckMate 816 trial. *Cancer Res*. 2021;81-(13 Supplement):CT003.
- Oezkan F, He K, Owen D, Pietrzak M, Cho JH, Kitzler R, et al. OA13.07 Neoadjuvant Atezolizumab in Resectable NSCLC patients: Immunophenotyping results from the interim analysis of the multicenter trial LCMC3. *Thorac Oncol*. 2019;14(10, Supplement):S242–S3.
- Cascone T, William WN Jr, Weissferdt A, Leung CH, Lin HY, Pataer A, et al. Neoadjuvant nivolumab or nivolumab plus ipilimumab in operable non-small cell lung cancer: the phase 2 randomized NEOSTAR trial. *Nat Med*. 2021;27(3):504–14.
- William WN Jr, Pataer A, Kalhor N, Correa AM, Rice DC, Wistuba II, et al. Computed tomography RECIST assessment of histopathologic response and prediction of survival in patients with resectable non-small-cell lung cancer after neoadjuvant chemotherapy. *J Thorac Oncol*. 2013;8(2):222–8.
- Lee HY, Lee HJ, Kim YT, Kang CH, Jang BG, Chung DH, et al. Value of combined interpretation of computed tomography response and positron emission tomography response for prediction of prognosis after neoadjuvant chemotherapy in non-small cell lung cancer. *J Thorac Oncol*. 2010;5(4):497–503.
- Poettgen C, Theegarten D, Eberhardt W, Levegruen S, Gauler T, Krbeik T, et al. Correlation of PET/CT findings and histopathology after neoadjuvant therapy in non-small cell lung cancer. *Oncology*. 2007;73(5–6):316–23.
- Forde PM, Spicer J, Lu S, Provencio M, Mitsudomi T, Awad MM, et al. Neoadjuvant Nivolumab plus chemotherapy in Resectable lung cancer. *N Engl J Med*. 2022;386(21):1973–85.
- Bar J, Urban D, Redinsky I, Ackerstein A, Daher S, Kamer I, et al. OA11.01 Neoadjuvant Pembrolizumab for early stage non-small cell lung cancer. *J Thorac Oncol*. 2021;16(10, Supplement):S865–S6.
- Xu C, Chen Q, Zhou C, Wu L, Li W, Zhang H, et al. 98P Camrelizumab as neoadjuvant, first- or later-line treatment for non-small cell lung cancer (NSCLC): a retrospective real-world study (CTONG2004). *Ann Oncol*. 2021;32:S1417.
- Travis WD, Dacic S, Wistuba I, Sholl L, Adusumilli P, Bubendorf L, et al. IASLC multidisciplinary recommendations for pathologic assessment of lung cancer resection specimens after Neoadjuvant therapy. *J Thorac Oncol*. 2020;15(5):709–40.
- Castello A, Rossi S, Lopci E. ^{18}F -FDG PET/CT in restaging and evaluation of response to therapy in lung cancer: state of the art. *Curr Radiopharm*. 2020;13(3):228–37.
- Zhong W-Z, Chen K-N, Chen C, Gu C-D, Wang J, Yang X-N, et al. Erlotinib versus gemcitabine plus cisplatin as neoadjuvant treatment of stage IIIA-N2 EGFR-mutant non-small-cell lung cancer (EMERGING-CTONG 1103): a randomized phase II study. *J Clin Oncol*. 2019;37(25):2235–45.
- Zhang C, Li S-L, Nie Q, Dong S, Shao Y, Yang X-n, et al. Neoadjuvant crizotinib in resectable locally advanced non-small cell lung cancer with ALK rearrangement. *J Thorac Oncol*. 2019;14(4):726–31.
- Kang J, Zhang C, Zhong WZ. Neoadjuvant immunotherapy for non-small cell lung cancer: state of the art. *Cancer Commun*. 2021;41(4):287–302.
- Lee J, Chaft J, Nicholas A, Patterson A, Waqar S, Tozola E, et al. PS01. 05 surgical and clinical outcomes with Neoadjuvant Atezolizumab in Resectable stage IB–IIIB NSCLC: LCMC3 trial primary analysis. *J Thorac Oncol*. 2021;16(3):S59–61.
- Aide N, Hicks RJ, Le Tourneau C, Lheureux S, Fanti S, Lopci E. FDG PET/CT for assessing tumour response to immunotherapy: report on the EANM symposium on immune modulation and recent review of the literature. *Eur J Nucl Med Mol Imaging*. 2019;46(1):238–50.
- Takada K, Toyokawa G, Okamoto T, Baba S, Kozuma Y, Matsubara T, et al. Metabolic characteristics of programmed cell death-ligand 1-expressing lung cancer on (18) F-fluorodeoxyglucose

- positron emission tomography/computed tomography. *Cancer Med.* 2017;6(11):2552–61.
23. Lopci E, Toschi L, Grizzi F, Rahal D, Olivari L, Castino GF, et al. Correlation of metabolic information on FDG-PET with tissue expression of immune markers in patients with non-small cell lung cancer (NSCLC) who are candidates for upfront surgery. *Eur J Nucl Med Mol Imaging.* 2016;43(11):1954–61.
 24. Tao X, Li N, Wu N, He J, Ying J, Gao S, et al. The efficiency of (18)F-FDG PET-CT for predicting the major pathologic response to the neoadjuvant PD-1 blockade in resectable non-small cell lung cancer. *Eur J Nucl Med Mol Imaging.* 2020;47(5):1209–19.

How to cite this article: Chen Z-Y, Fu R, Tan X-Y, Yan L-X, Tang W-F, Qiu Z-B, et al. Dynamic ^{18}F -FDG PET/CT can predict the major pathological response to neoadjuvant immunotherapy in non-small cell lung cancer. *Thorac Cancer.* 2022; 13(17):2524–31. <https://doi.org/10.1111/1759-7714.14562>